

## First Record of a Snailfish, *Careproctus notosaikaiensis* (Scorpaeniformes: Liparidae) from Korea

Hwan-Sung Ji<sup>1</sup>, Jeong-Ho Park<sup>2</sup>, Tae-Woo Ban<sup>1</sup>, Jin-Koo Kim<sup>1,\*</sup>

<sup>1</sup>Department of Marine Biology, Pukyong National University, Busan 608-737, Korea

<sup>2</sup>East Sea Fisheries Research Institute, NFRDI, Gangneung 210-861, Korea

### ABSTRACT

A single specimen (134.3 mm standard length) of a snailfish, *Careproctus notosaikaiensis* was collected from a fish trap in Goseong-gun, Gangwon-do, East Sea, Korea. It is characterized by having the teeth strongly trilobed; dorsal fin rays 52; anal fin rays 47; pectoral fin rays 35; caudal fin rays 10; vertebrae 58; ribs 2 pairs; cephalic pores, 2-6-7-2; gill slit extending to the fifth pectoral fin ray; chin pores paired and equal in size; dorsal and anal fins with distinct reddish margins. We describe this species as the first record to Korea, and proposed the new Korean name, “Dong-hae-bun-hong-ggom-chi” for this species.

**Keywords:** first record, snailfish, Liparidae, *Careproctus notosaikaiensis*, East Sea

### INTRODUCTION

The genus *Careproctus* Krøyer, 1862 belongs to the family Liparidae in the order Scorpaeniformes are distributed in Atlantic, Pacific, Antarctic, and Arctic waters, and are known mainly from the deeper waters of the continental slope (Chernova et al., 2004; Orr and Maslenikov, 2007). The genus comprises 112 species (Chernova et al., 2004; Orr and Maslenikov, 2007; Eschmyer and Fricke, 2011; Kai et al., 2011), of which 50 species are known from the North Pacific (Chernova et al., 2004), 22 from Japan (Nakabo, 2002; Sakurai and Shinohara, 2008; Kai et al., 2011), and a only single species, *Careproctus rastrinus*, from Korea (Kim et al., 2001, 2005).

In Korea, firstly *C. rastrinus* was reported by Kim et al. (1988) in the Fishes of the North Pacific Ocean; later, two species (*Careproctus furcellus*, *Careproctus* sp.) were described by Lee et al. (1999), also in the Fishes of the Pacific Ocean. The first description of *C. rastrinus* collected from Korea was made by Kim et al. (2001); later, Kim et al. (2005) reported the occurrence of *C. rastrinus* in Goseong and Sokcho, Gangwon-do, Korea.

*Careproctus* species are defined by the following combination of morphological characters: single pair of nostrils; presence of a pelvic disk; fewer pectoral fin rays than anal fin rays; absence of pseudobranchia; and non-variegated body color, except for the fins (Kido, 1988; Stein et al., 2001).

Recently, several taxonomic studies of *Careproctus* species have been conducted. Kido (1988) reported on the morphological characteristics of 18 *Careproctus* species; subsequently, some new species have been reported (Orr and Maslenikov, 2007; Sakurai and Shinohara, 2008; Kai et al., 2011).

A single specimen of the species *Careproctus notosaikaiensis* was collected from a fish trap in Goseong, Gangwon-do, East Sea, Korea. We here describe this first recorded instance of *C. notosaikaiensis* in Korea on the basis of this specimen. Counts and measurements followed those of Andriashev and Stein (1998). Cephalic pore terminology and the counts of pectoral fin rays followed those of Orr and Busby (2006). Each body part was measured to the nearest 0.1 mm using digital Vernier calipers. The numbers of vertebrae and fin rays were counted from radiographs (Hitex HA-100; Hitex Co., Tokyo, Japan). The present specimen of *C. notosaikaiensis* is deposited in the National Institute of Biological Resources (NIBR), Korea.

### SYSTEMATIC ACCOUNTS

Order Scorpaeniformes  
Family Liparidae Scopoli, 1777  
Genus *Careproctus* Krøyer, 1862

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*To whom correspondence should be addressed  
Tel: 82-51-629-5927, Fax: 82-51-629-5931  
E-mail: taengko@hanmail.net

<sup>1</sup>\**Careproctus notosaikaiensis* Kai, Ikeguchi, and Nakabo, 2011 (Table 1, Fig. 1)

*Careproctus notosaikaiensis* Kai, Ikeguchi, and Nakabo, 2011 (type locality: off Saikai, Japan).

**Material examined.** 1 specimen, 134.3 mm standard length (SL), Gangwon-do, Goseong-gun, 23 Aug 2011, Park JH, NIBR-P0000016325, from a fish trap at 200 m water depth.

**Description.** Dorsal fin rays 52; Anal fin rays 47; Pectoral fin rays 35; caudal fin rays 10; vertebrae 11+47=58; ribs, 2; gill rakers, 8; cephalic pores, 2-6-7-2.

Counts and proportional measurements are shown in Table 1. Body somewhat high, compressed, and tapering to tail. Head moderately large and slightly compressed; dorsal profile of head concave. Eye relatively large; pupil round. Single pair of tubular nostrils, located in front of eye. Snout rounded. Mouth subterminal; upper and lower jaw lengths equal; posterior tip of upper jaw reaching to middle part of eye. Teeth small, well-developed, and trilobed; arranged in 7–11 oblique rows clustered in wide bands on both jaws. Inner teeth strongly trilobed and larger than outer teeth. Cephalic pores well developed: nasal pores, 2; maxillary pores, 6; preoperculomandibular pores, 7; suprabranchial pores, 2. Chin pores paired and equal in size; present on ventral side of lower jaw. Upper margin of gill slit at level of dorsal margin of eye, extending to the 5th pectoral fin ray. Opercular flap angular and pointed slightly dorsally. Gill rakers short. Pectoral fin distinctly notched; upper lobe with 26 rays, not reaching origin of anal fin; lower lobe somewhat short, with 9 rays, extending beyond anus. Dorsal fin origin above middle of pectoral fin. Caudal fin narrow and truncate. Ventral disk small and rounded; anus close to disk.

**Color.** When fresh: body and head generally pinkish; abdominal cavity somewhat black; cephalic pores and tubular nostrils white; dorsal, anal, and caudal fins with distinct reddish margins; pectoral fin pale reddish; eye generally black without sclera. In ethanol: body and head pale white; abdominal cavity somewhat black; dorsal, anal, and caudal fins with black margins; pectoral and caudal fins pale white.

**Distribution.** Goseong, Gangwon-do, East Sea, Korea, at 200 m water depth (present study). Off Saikai, west coast of the Noto Peninsula, Japan (Kai et al., 2011).

**Remarks.** This specimen is identified as belonging to the genus *Careproctus*, based on the following characters: body generally pinkish; fewer pectoral fin rays than anal fin rays; absence of pseudobranchia; non-variegated body color, except for the fins (Kido, 1988; Stein et al., 2001). Based on comparisons with *Careproctus* species from the North Pacific, the specimen was identified as *C. notosaikaiensis* by numbers of dorsal, anal, pectoral, and caudal fin rays, and numbers of vertebrae, gill rakers, and ribs (Table 1). In particular, the specimen shows a strong correspondence with *C. notosaikaiensis*, in having: strongly trilobed teeth; paired, equal-sized chin pores; gill slits extending to the 5th pectoral fin ray; 2 pairs of ribs; distinctly reddish margins of the dorsal and anal fins (Kai et al., 2011). The specimen differs slightly from *C. notosaikaiensis* in the lengths of the caudal fins and preanal fins (Table 1); these differences are thought to represent regional variations, although genetic analyses are required for clarification. *Careproctus notosaikaiensis* is most similar to *C. rastrinus* in terms of meristic characters; however, they are easily distinguished by the shape of their teeth (trilobed in *C. notosaikaiensis* vs. conical in *C. rastrinus*) and the numbers of caudal fin rays (10 in *C. notosaikai-*



**Fig. 1.** *Careproctus notosaikaiensis*, NIBR-P0000016325, 134.3 mm standard length, collected from Goseong-gun, Gangwon-do.

Korean name: <sup>1</sup>\* 동해분홍꼼치

**Table 1.** Comparison of meristic and morphometric characters of *Careproctus notosaikaiensis*

	Present study (NIBR-P0000016325)	<i>Careproctus notosaikaiensis</i> Kai, Ikeguchi and Nakabo, 2011
No. of specimens	1	6
Standard length (SL, mm)	134.3	102.7–181.9
Measurements (% SL)		
Head length	28.0	27.6–31.4
Snout length	8.6	8.8–10.9
Eye diameter	5.6	5.5–6.4
Interorbital width	8.3	7.9–11.7
Body depth	27.6	24.1–29.5
Body width	14.7	12.8–15.7
Upper jaw length	11.4	11.1–12.5
Predorsal length	31.8	31.3–36.6
Preanal length	40.4	39.3–46.3
Preanus length	24.4	25.3–27.4
Pectoral fin length	17.1	15.3–19.3
Length of lower lobe of pectoral fin	14.5	13.3–20.4
Gill slit length	8.4	7.5–9.7
Disc length	5.8	5.6–7.0
Disc width	5.7	5.3–7.3
Caudal fin length	10.0	12.1–14.1
Counts		
Dorsal fin rays	52	52
Anal fin rays	47	46–47
Caudal fin rays	10	10
Pectoral fin rays	35	35–37
Total vertebrae	58	57–58
Gill rakers	8	7–11
Cephalic pores		
Nasal pores	2	2
Maxillary pores	6	6
Preoperculomandibular pores	7	7
Suprabranchial pores	2	2

*ensis* vs. 9 in *C. rastrinus*) (Nakabo, 2002; Kim et al., 2005). *Careproctus notosaikaiensis* is also similar to *Careproctus bowersianus* in terms of meristic characters, but they are distinguished by the numbers of ribs (2 in *C. notosaikaiensis* vs. 3 in *C. bowersianus*) and the colors of the dorsal and anal fins (distinctly reddish in *C. notosaikaiensis* vs. uniformly pinkish in *C. bowersianus*) (Gilbert and Burke, 1912; Burke, 1930). *Careproctus notosaikaiensis* is also similar to *Careproctus simus* in having strongly trilobed teeth, dark margined dorsal and anal fins and gill slits extending to pectoral fin ray; however, they are distinguished by the size of chin pores (equal size in *C. notosaikaiensis* vs. different size in *C. simus*) and the number of dorsal fin rays (52 in *C. notosaikaiensis* vs. 54–60 in *C. simus*) (Nakabo, 2002; Chernova et al., 2004). We here propose a new Korean name “Dong-hae-bun-hong-ggom-chi” for *C. notosaikaiensis*.

## ACKNOWLEDGMENTS

We thank Mr. Jin-Ho Park (Seoul National University, Korea)

for the collected of the species. This research was supported by the project on survey and excavation of Korean indigenous species of the National Institute of Biological Resources (NIBR) under the Ministry of Environment, Korea.

## REFERENCES

- Andriashev AP, Stein DL, 1998. Review of the snailfish genus *Careproctus* (Liparidae, Scorpaeniformes) in Antarctic and adjacent waters. *Contributions in Science*, 470:1-63.
- Burke V, 1930. Revision of the fishes of the family Liparidae. *Bulletin of the United States National Museum*, 150:1-204.
- Chernova NV, Stein DL, Andriashev AP, 2004. Family Liparidae Scopoli 1777: snailfishes. *California Academy of Sciences, Annotated Checklists of Fishes*, 31:1-72.
- Eschmyer WN, Fricke R, 2011. *Catalog of Fishes electronic version* [Internet]. California Academy of Sciences, Accessed 15 Mar 2012, <<http://researcharchive.calacademy.org/ichthyology/catalog/fishcat-main.asp/>>.
- Gilbert CH, Burke CV, 1912. Fishes from Bering Sea and Kamchatka. *Bulletin of the United States Bureau of Fisheries*,

- 30:31-96.
- Kai Y, Ikeguchi S, Nakabo T, 2011. A new species of the genus *Careproctus* (Liparidae) from the Sea of Japan. *Ichthyological Research*, 58:350-354.
- Kido K, 1988. Phylogeny of the family Liparidae, with the taxonomy of the species found around Japan. *Memoirs of the Faculty of Fisheries, Hokkaido University*, 35:1-125.
- Kim IS, Choi Y, Lee CL, Kim BJ, Kim JH, 2005. Illustrated book of Korean fishes. Kyo-Hak Publishing Co., Seoul, pp. 266-269 (in Korean).
- Kim YU, Gong Y, Lim JY, 1988. Classification of the commercial fishes in the Pacific ocean. *National Fisheries Research, Busan*, pp. 152-154 (in Korean).
- Kim YU, Myoung JG, Kim YS, Han KH, Kang CB, Kim JK, Ryu JH, 2001. Marine fishes of Korea. Hanguel Publishing Co., Busan, p. 221 (in Korean).
- Krøyer HN, 1862. Nogle Bidrag til Nordisk ichthyologi. *Naturhist Tidsskr*, 1:233-310.
- Lee JU, Kim YU, Park CY, Moon DY, Kim JB, Kim JK, Baik CI, Yang WS, Kim YS, Hwang SJ, An DH, Jo HS, Kim SS, Oh TY, Choi SG, Kim DN, Huh SH, Ryu JH, 1999. Fishes of the Pacific Ocean. Hanguel Publishing Co., Busan, pp. 317-320 (in Korean).
- Nakabo T, 2002. Liparidae. In: *Fishes of Japan with pictorial keys to the species*. English ed. (Ed., Nakabo T). Tokai University Press, Tokyo, pp. 665-677.
- Orr JW, Busby MS, 2006. Revision of the snailfish genus *Allocareproctus* Pitruk & Fedorov (Teleostei: Liparidae), with descriptions of four new species from the Aleutian Islands. *Zootaxa*, 1173:1-37.
- Orr JW, Maslenikov KP, 2007. Two new variegated snailfishes of the genus *Careproctus* (Teleostei: Scorpaeniformes: Liparidae) from the Aleutian Islands, Alaska. *Copeia*, 2007: 699-710.
- Sakurai H, Shinohara G, 2008. *Careproctus rotundifrons*, a new snailfish (Scorpaeniformes: Liparidae) from Japan. *Bulletin of the National Museum of Nature and Science, Series A, Supplement*, 2:39-45.
- Scopoli GA, 1777. *Introductio ad historiam naturalem, sistens genera lapidum, plantarum et animalium hactenus detecta, characteribus essentialibus donata, in tribus divisa, subinde ad leges naturae*. Wolfgangum Gerle, Prague, pp. 1-506.
- Stein DL, Chernova NV, Andriashev AP, 2001. Snailfishes (Pices: Liparidae) of Australia, including descriptions of thirty new species. *Records of the Australian Museum*, 53: 341-406.

Received May 24, 2012  
 Revised September 3, 2012  
 Accepted September 3, 2012